

Constants & Units

Physical Constants

a_0	Bohr radius	5.29177×10^{-9} cm
c	speed of light in vacuum	2.99792×10^{10} cm s ⁻¹
e	electron charge	-4.80325×10^{-10} statcoulomb (= esu)
eV	electron volt	1.60218×10^{-12} erg
G	gravitational constant	6.67428×10^{-8} dyne cm ² g ⁻²
h	Planck's constant	6.62607×10^{-27} erg s
k	Boltzmann's constant	1.38065×10^{-16} erg K ⁻¹
m_e	electron mass	9.10938×10^{-28} g
m_p	proton mass	1.67262×10^{-24} g
R_∞	Rydberg constant	1.09737×10^5 cm ⁻¹
$R_\infty c$	Rydberg frequency	3.28984×10^{15} s ⁻¹
σ	Stefan – Boltzmann constant	5.67040×10^{-5} erg cm ⁻² s ⁻¹ K ⁻⁴
σ_T	Thomson cross section	6.65245×10^{-25} cm ²
u	atomic mass unit	1.66054×10^{-24} g

Astronomical Constants

au	astronomical unit	1.49598×10^{13} cm
H_0	Hubble constant	72 km s ⁻¹ Mpc ⁻¹
kpc	kiloparsec	10^3 pc
L_\odot	solar bolometric luminosity	3.826×10^{33} erg s ⁻¹
ly	light year	9.4605×10^{17} cm
M_\odot	solar mass	1.989×10^{33} g
Mpc	megaparsec	10^6 pc
pc	parsec	3.0856×10^{18} cm
R_\odot	solar radius	6.9598×10^{10} cm
yr	year	$3.156 \times 10^7 \approx 10^{7.5}$ s

MKS and Gaussian CGS Units

Type	mks unit	cgs unit	conversion
mass	kg	g	10^3 g = 1 kg
length	m	cm	10^2 cm = 1 m
time	s	s	

frequency	Hz	Hz	$1 \text{ Hz} = 1 \text{ s}^{-1}$
charge	coulomb	statcoulomb	$3 \times 10^9 \text{ statcoulomb} = 1 \text{ coulomb}$ (1 statcoulomb = 1 esu)
current	ampere	statampere	$3 \times 10^9 \text{ statamp} = 1 \text{ amp}$ (1 amp = 1 coulomb s ⁻¹)
electric field	v m ⁻¹	statvolt cm ⁻¹	$(1/3) \times 10^{-4} \text{ statvolt cm}^{-1} = 1 \text{ v m}^{-1}$
energy	joule	erg	$10^7 \text{ erg} = 1 \text{ joule}$
force	newton	dyne	$10^5 \text{ dyne} = 1 \text{ newton}$
magnetic field	tesla	gauss	$10^4 \text{ gauss} = 1 \text{ tesla}$
resistance	Ohm	sec cm ⁻¹	$(1/9) \times 10^{-11} \text{ s cm}^{-1} = 1 \text{ Ohm}$
temperature	Kelvin	Kelvin	

Other Constants and Units

arcmin	1/60 deg
arcsec	1/60 arcmin
Angstrom	10^{-10} m
dB	$0.1 \log_{10}(P_1/P_2)$
e	2.71828...
GHz	10^9 Hz
Jy	$10^{-26} \text{ W m}^{-2} \text{ Hz}^{-1}$ (1 mJy = 10^{-3} Jy, 1 μ Jy = 10^{-6} Jy)
MHz	10^6 Hz
μm	10^{-6} m
π	3.14159...
radian	$(180/\pi) \text{ deg} \approx 206264.8 \text{ arcsec}$

Engineers and physicists prefer mks (meter, kilogram, second) units, so most radio astronomers use mks units to describe their equipment and the results of their observations. Most astrophysicists prefer Gaussian cgs (centimeter, gram, second) units to describe astronomical sources. Thus you have to deal with both systems of units and be able to convert between them in order to do astronomy. J. D. Jackson's *Classical Electrodynamics* has an appendix explaining the different systems in detail.